



Research Article

August 2017 | Volume 5 | Issue 2 | Pages 180-185

ARTICLE INFO

Received
March 07, 2017
Accepted
May 19, 2017
Published
August 15, 2017

***Corresponding Author**

Kiplimo Jepkorir Joyce

E-mail

jkiplimo@kabianga.ac.ke

Keywords

Antibacterial activity
Conyza bonariensis
Essential oil composition
Hydrocarbons
Therapeutic properties

How to Cite

Musembei R, Joyce KJ.
Chemical composition and
antibacterial activity of
essential oil from Kenyan
Conyza bonariensis (L.)
Cronquist. Sci Lett 2017;
5(2):180-185



Scan QR code to see this
publication on your
mobile device.

Open Access

Chemical Composition and Antibacterial Activity of Essential Oil from Kenyan *Conyza bonariensis* (L.) Cronquist

Racheal Musembei, Kiplimo Jepkorir Joyce*

Department of Physical Sciences, School of Science, University of Kabianga, 20200, Kericho, Kenya

Abstract

The utilization of plants to treat sicknesses and to prevent diseases, including epidemics, has been in practice for thousands of years. The understanding of their therapeutic properties has been passed over the centuries within and among individual communities. Secondary metabolism produces active compounds which are usually responsible for the biological and pharmacological properties of some plant species used for the management of communicable diseases. Presently, the antimicrobial activity of several plants, up to now considered empirical has been scientifically established. Several studies have aimed to explain the chemical composition and the mechanisms involved in microbial growth inhibition, either separately or together with conventional antimicrobial compounds. In this study, the essential oil from aerial parts of *Conyza bonariensis* (L.) Cronquist collected from Kabianga division, Kericho County, Kenya was obtained by hydro-distillation and analyzed by GC-MS. The major components were monoterpene hydrocarbons; limonene (8.26%), *trans*-ocimene (3.74%) and sesquiterpene hydrocarbon *trans*- β -farnesene (3.17%), and among other compounds isolated an aldehyde, 1H-Indene-3-carboxaldehyde, 2,6,7,7a-tetrahydro-1,5-dimethyl yielded 49.14%. The antibacterial activity indicated that the oil was active against the bacterial strains tested. The minimum inhibitory concentration for the oil against *Escherichia coli* was 12.5%, while that of *Salmonella typhi* was 6.25%. The results suggested that the essential oil of *C. bonariensis* can be developed as a safe, natural and alternative method for controlling bacterial infections.



This work is licensed under the Creative Commons Attribution-Non Commercial 4.0 International License.

Introduction

Plants and their essential oils are potentially useful sources of antimicrobial compounds. There are numerous studies and published data on the antimicrobial activities of plant compounds against many different types of microbes. The main constituents of essential oils mono- and sesquiterpenes including carbohydrates, phenols, alcohols, ethers, aldehydes and ketones are responsible for the biological activity of aromatic and medicinal plants as well as for their fragrance. Due to these properties, spices and herbs have been added to food since ancient time, not only as flavoring agents but also as preservatives [1]. The genus *Conyza* belongs to the family *Asteraceae* and comprises about seventeen species of annual, biennial and perennial herbs, particularly inhabiting highland and mountain regions [2]. Seven species have been reported: *Conyza canadensis* (L.) Cronquist, *C. laevigata* (Rick) Pruski, *C. primulifolia* (Lam.) Cuatrec. & Lourteig, *C. sumatrensis* (Retz) E. Walker, *C. uliginosa* var. *columbiana* (Hieron) Cuatrec, *C. uliginosa* var. *uliginosa* (Benth) Cuatrec and *C. bonariensis* (L) Cronquist [3]. *C. bonariensis* belongs to the flowering plants; it grows widely in Kenya, and is known as ‘taplile’ by the Kalenjini community of Kenya. An infusion from *C. bonariensis* aerial parts is used in traditional medicines as antiseptic, hemostatic and antispasmodic [4]. The leaves are also used as a decoction in the treatment of malaria, fever, jaundice, skin rashes, sickle cell anemia, inflammations of the fallopian tubes and backache [5]. In women, the leaves are used to induce uterine contraction, abortion and stimulate lactation [6]. Diseases such as rheumatism, cystitis, gout, nephritis, toothache, and headache have been managed by the use of *C. bonariensis* [5]. Some of the biological activities of the *Conyza* genus are: *C. sumatrensis*, antimalarial activity [4]; *C. Canadensis*, antibacterial, antioxidant, cytotoxic and anti-proliferative activity [7]; *C. aegyptiaca*, anti-inflammatory, analgesic and antiviral activity [8-10]; *C. dioscorides*, antischistosomal activity [11]; *C. filaginoides*, anti-protozoa and anti-diarrheal activity [12, 13] and *C. bonariensis*,

molluscicidal, anti-inflammatory, antipyretic and antimicrobial properties [14]. Previous investigations of *C. bonariensis* have revealed glycosides, polyphenolic compounds, flavonoids sesquiterpenic lactones and essential oils [15, 16]. The current research was done to look at the chemical constituents and antibacterial activity of the essential oil from the Kenyan *C. bonariensis*; this is the first report on the essential oil of the Kenyan *Conyza* species.

Materials and Methods

Plant material

C. bonariensis (L.) Cronquist leaves were collected in Kabianga location, Kericho, Kenya, in February 2016, and identified by a botanist from the Biological Sciences Department. A voucher specimen (Cb 01) was deposited in the University herbarium, School of Agriculture and Biotechnology, Kabianga.

Essential oil isolation

Fresh leaves (about 1000 g) were subjected to hydro-distillation for 4 h using a Clevenger-type apparatus (Garg process glass, India). This apparatus consists of three parts: a round bottomed flask in which the material containing essential oil and given the quantity of water are placed, a separator in which the oil is automatically separated from the distillate and a condenser. The oil (0.4 ml) was dried over anhydrous sodium sulfate in order to remove any traces of water and then stored in sealed vials at 2 °C in the dark until analyzed and tested. The yield (0.04%) was calculated based on the dry weight of the plant material.

Gas chromatography-mass spectrometry

About 1 mg of *C. bonariensis* essential oil was diluted in 1 ml volume of dichloromethane and analyzed by gas chromatography–mass spectroscopy (GC-MS). The peak area was used for relative quantification. The essential oil was analyzed in full scan mode of GC-MS. A GC column HP-5 MS ultra-inert column (30 m × 0.25 mm; 0.25 μm) (J & W, Folsom, CA, USA) with the flow rate of 1.2 ml/min was used. The injection mode was splitless and the oven temperature of 35 °C was maintained

for 5 min and then programmed to 280 °C at 10°C/min for 10.5 min. The total run time was 50 min and the injection volume was 1 µl. The mass spectrometer (MS) was operated in the positive electron impact (EI) mode at 70 eV, in m/z range of 42-350. The identification of the compounds was done by comparing their retention times and mass spectra with those found in NIST 11 chemical mass spectral database. The relative proportions of the essential oil constituents were expressed as a percentage obtained by the peak area normalization.

Antibacterial assay

The antibacterial potential of the essential oil was tested according to the national committee of clinical laboratory (NCCL) standards at the Biotechnology Laboratory, Egerton University. The Gram-negative bacterial strains *Escherichia coli* ATCC 25922 and *Salmonella typhi* ATCC 13311 were used for the analysis. The antibacterial activity was determined using disc diffusion method [17]. All the apparatus for the culturing process were sterilized at 120°C for 20 minutes. The Mueller Hinton agar was used as the plating media to culture both bacterial strains [18]. The bacterial strains of *E. coli* and *S. typhi* were primarily incubated in the nutrient broth at a temperature of 37°C to 0.5 on the McFarland scale. The cultures were then diluted to the concentration of 1.0×10^5 CFU/ml with sterilized water, transferred to Muller Hilton media and added into the Petri plates [19]. The stock solution weighed 1 ml of the pure oil was serially diluted to 75%, 50%, 25%, 12.5%, and 6.25 % using dimethyl sulfoxide as a diluting agent. Later, 4 mm filter paper discs were placed on the medium and added with different dilutions of *C. bonariensis* oil in triplicate. The pure dimethyl sulfoxide was used as a control. Later, the plates were incubated at 37°C and the inhibition zones were measured in mm after two days [20].

Results and discussion

The hydrodistilled essential oil from the leaves of *C. bonariensis* was analyzed by GC-MS and 37 compounds were identified (Fig. 1). The retention times and area percentages of the compounds identified by GC-MS are given in Table 1. The

Table 1 The chemical composition of *C. bonariensis* essential oils as determined by gas chromatography-mass spectroscopy.

RT (min)	Compound name	Percentage
Monoterpene hydrocarbons		
9.48	α-Phellandrene	0.90
9.58	α- Pinene	2.25
10.21	Benzaldehyde	2.20
10.52	β-Pinene	5.37
10.84	Myrcene	1.30
11.48	o-Cymene	0.82
11.56	Limonene	8.26
11.73	Sylvestrene	0.90
11.92	(E)-β-Ocimene	3.74
12.11	γ- Terpinene	0.93
12.63	Terpinolene	0.70
13.29	neo-allo-Ocimene	0.70
13.33	Verbenene	0.70
14.11	Terpinen-4-ol	0.85
16.46	Camphene	1.10
Oxygenated monoterpenes		
16.73	Eugenol	0.68
Sesquiterpene hydrocarbons		
17.00	α-Copaene	0.68
17.06	Daucene	0.89
17.19	β-Elementene	0.86
17.61	(E)-Caryophyllene	1.59
17.73	β-Cubebene	1.14
17.96	(E)-β-farnesene	3.17
18.05	α-Humulene	1.01
18.16	trans-Muurola-4(14),5-diene	0.79
18.27	γ- Cadinene	1.47
18.39	Germacrene D	1.55
18.58	Bicyclogermacrene	1.02
20.08	γ- Muurolene	0.81
Oxygenated sesquiterpenes		
19.39	Nerolidol	1.47
19.66	Spathulenol	1.32
19.82	1,10-Cubeno	1.65
Others		
18.78	1H-Indene-3-carboxaldehyde, 2,6,7,7a-tetrahydro-1,5-dimethyl-	49.14

RT = retention time

results indicated that, the dominant constituent in the oil was the monoterpene hydrocarbons, which accounted for 30.72%, of which limonene (8.26%), β- pinene (5.37 %) and β-ocimene (3.74%) were the major compounds. The previous investigation of the leaf oil of *C. bonariensis* collected from Merida, Venezuela also showed that these monoterpenes were the main compounds; however, a higher concentration of trans-β-ocimene (20.7%) was also reported [20]. The sesquiterpene hydrocarbons accounted for 14.98% of the total oil, with β-farnesene (3.17%), E-caryophyllene (1.59%) and γ-cadinene (1.47%) being the major compounds. These results were in accordance with those of Lambert et al. [17] who found that trans-β-farnesene




Fig. 1 Representative total ion chromatogram of *Conyza bonariensis* essential oils with retention time of each compound detected by gas chromatography–mass spectroscopy.

(37.8%), trans-ocimene (20.7%) and β -sesquiphellandrene (9.8%) were present as the major components of the *C. bonariensis* collected from Venezuela. The findings of this study were also in agreement with those of Mabrouk et al. [16] and Maia et al. [21], who also reported *trans*- β -farnesene as one of the main compounds of the essential oil of this species. On the other hand, Šoškić et al. [22] reported limonene and *trans*-ocimene as the main proportions in the essential oil of *C. bonariensis* found in Rio de Janeiro, Brazil. β -Sesquiphellandrene was only observed in low amounts in the essential oil studied by Souza et al. [23]. Limonene was detected in 5.1% in the same species collected in Rio de Janeiro, Brazil [24]. The oxygenated derivatives of both monoterpenes and sesquiterpene contributed to 0.68% and 4.44%, respectively, it was found that nerolidol, spathulenol and 1,10-cubeno were the major components in the present investigation. *C. bonariensis* found in Kenya and reported herein had the same components as those found in other regions but in low concentration. This variability in the chemical

composition could be attributed to numbers of factors such as genetics (species or variety) of the plant, environmental conditions (geographical, climatic or seasonal) and processing of plant material. Variability in the qualitative and quantitative composition most probably depended on the genotype of the plant and the influence of different environmental factors [22].

The essential oil was tested for antibacterial activity against the pathogenic Gram-negative bacterial strains (*E. coli* and *S. typhi*). The oil exhibited activity against the two bacterial strains tested as seen by their zones of inhibition (Table 2). However, the activity of the oil is known to differ with its concentration and the kind of bacteria. The activity response to *S. typhi* was greater than in *E. coli* at different concentrations, except 50% oil concentration. The minimum inhibitory concentration for the oil against both bacterial strains was 12.5% for *E. coli* and 6.25% for *S. typhi*. The essential oil tested in this work has a range of compounds that could account for antibacterial activity as a result of their synergistic effect.

Table 2 The antibacterial activity of different concentration of essential oil of *Conyza bonariensis* against *Escherichia coli* and *Salmonella typhi*.

Microorganisms	Zone of Inhibition (mm)					
	100%	75%	50%	25%	12.5%	6.25%
<i>Escherichia coli</i> ATCC 25922	14.7 ± 1.3	11 ± 0.8	8.7 ± 0.5	7.7 ± 0.4	6.3 ± 0.5	0.0 ± 0.0
<i>Salmonella typhi</i> ATCC 13311	16 ± 0.0	12 ± 0.8	8.7 ± 0.5	7.3 ± 0.5	7.3 ± 1.3	6.0 ± 0.2

The essential oil was diluted with dimethyl sulfoxide and antibacterial activity was determined using disc diffusion assay.

Although, essential oils usually occur as complex mixtures of compounds and its antimicrobial activity can generally be accounted for their major monoterpene components. This is because that the monoterpenes can diffuse into and harm cell membrane structures [24]. Previous research indicated that, limonene, β -pinene, β -ocimene, β -caryophyllene and β -farnesene possess moderate antibacterial activity [25], it has also been reported that oxygenated components of essential oil have a higher potential, mainly phenol-type compounds, for example, thymol and carvacrol while hydrocarbon monoterpenes demonstrate the lowest antibacterial activity. Other reports indicated that oxygenated monoterpenes, display strong antimicrobial activity, mostly on whole cells, while hydrocarbon compounds show lower activity, because of their limited diffusion through the medium insolubility in water [26]. Generally, hydrocarbons are relatively inactive regardless of their structural type, and this inactivity is closely related to their limited hydrogen bond capacity and water solubility [27]. Ketones, aldehydes and alcohols are active, but with differing specificity and levels of activity, which is related to the present functional group, but also associated with hydrogen-bonding parameters in all cases. Previous results showed that greater antimicrobial potential could be ascribed to the oxygenated terpenes, especially phenolic compounds [28].

Conclusions

In this study, the essential oil from the plant *C. bonariensis* was extracted and analyzed. The results showed that oil obtained could be developed into a natural antibiotic because its compounds demonstrated potential activity against bacterial strains. These components displayed potential for synergistic coupling with antimicrobial agents to advance therapeutic effectiveness, in the face of increasing bacterial resistance; however, this needs

further investigation.

Acknowledgements

The authors wish to thank Mr. Hillary Kirwa of International Centre of Insect Physiology and Ecology (ICIPE) center in Kenya for GC-MS analyses.

Conflict of interest

There is no conflict of interest.

References

- [1] Kalim M, Hussain F, Ali H, Iqbal MN. Antifungal activities of methanolic extracts of *Datura innoxia*. PSM Biol Res 2016; 1(2):70-73.
- [2] Buckingham J, Macdonald FM, Bradley HM. Dictionary of Natural Products. Type of Compounds Index. Chapman and Hall, London UK; 1994.
- [3] Badillo VM. Lista actualizada de las especies de la familia Compuesta (*Asteraceae*) de Venezuela. Ernstia, 2001; 11:147-215.
- [4] Boniface PK, Pal A. Substantiation of the Ethnopharmacological use of *Conyza sumatrensis* (Retz) E.H. Walker in the treatment of malaria through *in-vivo* evaluation in *Plasmodium berghei* infected mice. J Ethnopharmacol. 2013; 145:373-377.
- [5] Cheng XR, Thabit RA, Wang W, Shi HW, Shi YH, Le GW. Analysis and comparison of the essential oil in *Conyza bonariensis* grown in Yemen and China. Prog Biomed, 2013; 36:011.
- [6] Okello SV, Nyunja RO, Netondo GW, Onyango JC. Ethnobotanical study of medicinal plants used by Sabaots of Mt. Elgon Kenya. Afr J Tradit Complement Altern Med 2010; 7:1-10.
- [7] Hayet E, Maha M, Samia A, Mahjoub MA, Brahim S, Kenani A, Zine M, Aouni M. Antibacterial, antioxidant and cytotoxic activities of extracts of *Conyza canadensis* (L.) Cronquist growing in Tunisia. Med Chem Res 2009; 18:447-454.
- [8] Heras B, Slowing K, Benedi, J, Carretero E, Ortega T, Toledo C, Bermejo P, Iglesias I, Abad MJ, Gómez-Serranillos P, Liso PA, Villar A, Chiriboga X. Antiinflammatory and antioxidant activity of plants used in traditional medicine in Ecuador. J Ethnopharmacol 1998; 61:161-166.
- [9] Asogalem EA, Foyet HS, Ngogang J, Folefoc GN, Dimo T, Kamtchoung P. Analgesic and anti-inflammatory activities of *Erigeron floribundus*. J Ethnopharmacol 2004; 91:301-308.
- [10] Anani K, Hudson JB, De Sousa C, Akpagana K, Tower GH, Arnason JT, Gbeassor M. Investigation of medicinal plants of Togo for antiviral and antimicrobial activities. Pharm Biol 2000;

- 38:40-45.
- [11] Kamel EG, El-Emam MA, Mahmoud SS, Fouda FM, Bayaomy FE. Parasitological and biochemical parameters in *Schistosoma mansoni*-infected mice treated with methanol extract from the plants *Chenopodium ambrosioides*, *Conyza dioscorides* and *Sesbania sesban*. Parasitol Intern 2011; 60:388-392.
- [12] Calzada F, Cedillo-Rivera R, Mata R. Antiprotozoal activity of the constituents of *Conyza filaginoides*. J Nat Prod 2001; 64:671-673
- [13] Atta AH, Mounier, SM. Antidiarrhoeal activity of some Egyptian plant extracts. J Ethnopharmacol 2004; 92: 303-309.
- [14] Thabit RAS, Cheng X, Al-Hajj N, Rahman MRT, Lei, G. Antioxidant and *Conyza bonariensis*: A review. Eur Acad Res 2014; 2:8454-8474.
- [15] Zahoor A, Siddiqui IN, Khan A, Ahmad VU, Ahmed A, Hassan Z, Khan S, Iqbal S. Two new glycosides from *Conyza bonariensis*. Nat Prod Comm 2010; 5:1099-1102.
- [16] Hussein K, Ahmed AH, Al-Maqtari MA. Composition and radical scavenging activity of edible wild *Pulicaria jaubertii* (Asteraceae) volatile oil. PSM Biol Res 2017; 2(1):21-29.
- [17] Lambert RJ, Skandamis PN, Coote PJ, Nychas GJ. A study of the minimum inhibitory concentration and mode of action of oregano essential oil, thymol and carvacrol. J App Microbiol 2001; 91:453-462.
- [18] Mueller JH, Hinton J. A protein-free medium for primary isolation of gonococcus and meningococcus. Proc Soc Exp Biol Med 1941; 48:3330-333.
- [19] Araujo L, Moujir LM, Rojas J, Rojas L, Carmona J, Rondón M. Chemical Composition and Biological Activity of *Conyza Bonariensis* Essential Oil Collected in Mérida, Venezuela. Nat Prod Comm 2013; 8:1-4.
- [20] Daouk KD, Dagher MS, Sattout JE. Antifungal activity of the essential oil of *Origanum syriacum* L. J Food Prot 1995; 58:1147-1149.
- [21] Maia JGS, da Silva MHL, Zoghbi MGB, Andrade EHA. Composition of the essential oils of *Conyza bonariensis* (L.) Cronquist. J Essent Oil Res 2002; 14:325-326.
- [22] Šoškić M, Bojović D, Tadić V. Comparative chemical analysis of essential oils from Lavender of different geographic origins. Studia Ubb Chemia, Lxi 2016; 2:127-136.
- [23] Souza MC, Siani AC, Ramos MF, Menezes-de-Lima O, Henriques MGMO. Evaluation of anti-inflammatory activity of essential oils from two *Asteraceae* species. Pharmazie 2003; 58:582-586.
- [24] Sikkema J, de Bont JAM, Poolman B. Mechanisms of membrane toxicity of hydrocarbons. Microbiol Rev 1995; 59:201-222.
- [25] Soković M, Glamočlija J, Marin PD, Brkić D, van Griensven LJLD. Antibacterial effects of the essential oils of commonly consumed medicinal herbs using an *in vitro* model. Molecules 2010; 15:7532-7546.
- [26] Knobloch K, Weigand, H, Weis N, Schwarm HM, Vigneschow H. Action of terpenoids on energy metabolism. In: Progress in Essential Oil Research; Brunke, E.J., Ed.; Walter de Gruyter: Berlin, Germany; 1986, p. 429-445.
- [27] Griffin GS, Markham LJ, Leach ND. An agar dilution method for the determination of the minimum inhibitory concentration of essential oils. J Essent oil Res 2000; 12:149-255.
- [28] Soković M, Tzakou O, Pitarokili D, Couladis M. Antifungal activities of selected aromatic plants growing wild in Greece. Nahrung 2002; 46:317-320.